

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification</b> <sup>6</sup> : <b>A61K 9/50, 31/60</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 98/26767</b> <b>(43) International Publication Date:</b> 25 June 1998 (25.06.98)
<b>(21) International Application Number:</b> PCT/IB97/01652 <b>(22) International Filing Date:</b> 16 December 1997 (16.12.97) <b>(30) Priority Data:</b> 08/767,980 17 December 1996 (17.12.96) US <b>(71) Applicant (for all designated States except US):</b> POLI INDUSTRIA CHIMICA S.P.A. [IT/IT]; Piazza Agrippa, 1, I-20141 Milano (IT). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> Busettil, Cesare [IT/IT]; Via Signorini, 7, I-20149 Milano (IT). CRIMELLA, Tiziano [IT/IT]; Via Londonio, 1, I-20154 Milano (IT). OLGIATI, Vincenzo [IT/IT]; Via Settembrini, 16, I-20010 Casorezzo (IT). <b>(74) Agents:</b> McCallum, William, Potter et al.; Cruikshank & Fairweather, 19 Royal Exchange Square, Glasgow G1 3AE (GB).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> SITE-SPECIFIC CONTROLLED RELEASE DOSAGE FORMULATION FOR MESALAMINE  <b>(57) Abstract</b> <p>Pharmaceutical formulation for the site-specific delivery of mesalamine in the colon is disclosed. The formulation includes (a) a core comprising mesalamine in an amount effective to produce a therapeutic anti-inflammatory effect; (b) a swellable polymeric coating layer substantially surrounding the core which inhibits the release of mesalamine for a predetermined period of time dependent upon the thickness of the swellable polymeric coating layer, and (c) an outer enteric coating layer substantially surrounding the swellable polymeric coating layer. Methods of achieving the site-specific delivery of mesalamine in the colon are also disclosed.</p> <p style="text-align: center; font-size: 2em; font-family: cursive;">Hydrophilic matrix</p>		

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## **SITE-SPECIFIC CONTROLLED RELEASE DOSAGE FORMULATION FOR MESALAMINE**

### **Field of the Invention**

The present invention relates to controlled release dosage formulations. More particularly, the present invention relates to site-specific controlled release dosage formulations.

5

### **Background of the Invention**

Mesalamine, also commonly known as "mesalazine," "5-amino salicylic acid", and "5-ASA", is well known for its anti-inflammatory properties. It is classified a non-steroidal anti-inflammatory drug (N-SAID). Mesalamine has been approved for the treatment of mildly to moderately active ulcerative colitis.

10

With regard to the action of mesalamine in the treatment of ulcerative colitis and other inflammatory bowel diseases, there exists clear

-2-

evidence that i) the effect of mesalamine for treatment of inflammatory bowel diseases is exerted through a local rather than systemic action; and ii) the larger the systemic absorption the higher the possibility of side-effects (including renal necrosis, and pyuria). It is therefore advantageous to achieve maximum mesalamine concentration at the site of the disease, with the smallest possible systemic absorption. Since early release in the small bowel increases the systemic absorption and reduces the drug available for the topical action in the colon, the most powerful system is a technology which completely releases the active principle in the colon.

For this reason, attempts have been made to formulate mesalamine in a manner which will permit the delivery of the drug in the colon, i.e., at the site of the disease, to maximize the therapeutic effectiveness of the drug.

Delivery directly in the colon requires formulations which are capable of passing over the entire tract of the small intestine, including the duodenum, jejunum, and ileum, so that the active ingredients are released directly in the colon. Such formulations typically employ coatings for the purpose of preserving the integrity of the formulation while passing through the gastric tract. The high acidity, and presence of proteolytic and other enzymes generates a highly digestive environment which readily dissolves pharmaceutical formulations which do not possess some type of gastro-resistance protection.

Several formulations of mesalamine for delivery in the colon have been proposed. One approved mesalamine formulation is commercially available from Proctor & Gamble Pharmaceuticals under the trade name ASACOL® delayed release tablets. The formulation includes a tablet of mesalamine coated with the commercially available enteric coating EUDRAGIT®-S, which is a methacrylic acid copolymer which dissolves at pH 7 or greater.

Another formulation for mesalamine is commercially available from Marion Merrell Dow under the trade name PENTASA®, based on U.S. Patent Nos. 4,496,553 and 4,908,173 to both Halskov. The formulation includes a tablet formed from a granulate of mesalamine and

-3-

polyvinylpyrrolidone prepared using an organic solvent, which granulate is coated prior to the formation of the tablet, with a cellulose derivative coating material that will gradually release the active.

Solvay provides two formulations of mesalamine under the trade name ROWASA®. The formulations include a rectal suppository formulation and a rectal suspension enema.

In addition to the foregoing formulations, European Patent No. 366,621 describes a formulation for selective colon delivery of various drugs including ketoprofen and ibuprofen. The formulation includes a core coated with three different layers: an inner layer including an anionic polymer, an outer gastro-resistant layer and an intermediate swellable layer constituted by high viscosity cellulose derivatives of high molecular weight. Drug release in this system is dependent upon the pH to which the formulation is exposed. The inner layer is a polymer which is soluble only at a pH value of 7 or higher, and this layer must be dissolved before the release of mesalamine can begin.

Accordingly, there remains a continuing need in the art for pharmaceutical formulations which release active ingredient at a predetermined site or location in the body after a predetermined latency or lag time period, i.e., site-specific release formulations. In particular, there remains a need in the art for a pharmaceutical formulation for the site-specific delivery of mesalamine which achieves effective release of mesalamine directly into the colon. There further remains a need in the art for a method of preparing a pharmaceutical formulation for the site-specific delivery of mesalamine which does not require the use of organic solvents, or very dilute solutions of coating materials.

### Summary of the Invention

It is therefore an object of the present invention to provide a pharmaceutical formulation for the site-specific delivery of mesalamine. It is a further object of the present invention to provide a site-specific, controlled

-4-

release pharmaceutical formulation for mesalamine which provides release of mesalamine from the formulation at the predetermined site of delivery.

As a first aspect, the present invention provides a pharmaceutical formulation for the site-specific delivery of mesalamine in the colon. The formulation comprises: *a*) a core comprising mesalamine in an amount effective to produce a therapeutic anti-inflammatory effect; *b*) a swellable polymeric coating layer substantially surrounding the core which inhibits the release of mesalamine for a predetermined period of time dependent upon the thickness of the swellable polymeric coating layer; and *c*) an outer enteric coating layer substantially surrounding the swellable polymeric coating layer. The outer enteric coating layer inhibits the swelling of the swellable, intermediate coating layer until the formulation reaches the duodenum and dissolves upon exposure to pH above 4.5. The dissolution, erosion or disintegration of the outer enteric coating layer triggers the subsequent swelling and dissolution of the swellable polymeric coating layer.

As a second aspect, the present invention provides a method for achieving the site-specific delivery of mesalamine in the colon of a subject in need of such treatment. The method comprises orally administering to a subject in need thereof, a site-specific dosage formulation including: *a*) a core comprising mesalamine in an amount effective to produce a therapeutic anti-inflammatory effect; *b*) a swellable polymeric coating layer substantially surrounding the core, wherein the swellable polymeric coating layer inhibits the release of mesalamine for a predetermined period of time dependent upon the thickness of the swellable polymeric coating layer; and *c*) an outer enteric coating layer substantially surrounding the swellable polymeric coating layer, wherein the outer enteric coating dissolves upon exposure to pH greater than about 4.5, and wherein the dissolution of the outer enteric coating layer initiates the swelling and dissolution of the swellable polymeric coating layer.

-5-

The foregoing and other objects and aspects of the present invention are explained in detail in the detailed description and examples set forth hereinbelow.

### Detailed Description of the Invention

5           The pharmaceutical formulations and methods of the present invention provide a site-specific controlled-release pharmaceutical formulation for mesalamine and other nonsteroidal anti-inflammatory drugs which are desirously delivered in a site-specific manner to the colon.

10           Generally, the pharmaceutical formulations of the present invention include a core, a swellable polymeric coating layer and an outer enteric coating layer which dissolves upon exposure to a pH greater than about 4.5.

          The core is comprised of the active ingredient, i.e., mesalamine. In addition, the core typically also includes one or more pharmaceutically acceptable excipients. Pharmaceutically acceptable excipients which may be  
15           employed are well known to those skilled in the art and include any conventional pharmaceutically acceptable tableting excipients. Examples of suitable excipients which may be included in the core of the formulations of the present invention include but are not limited to microcrystalline cellulose, dibasic calcium phosphate dihydrate, starch, magnesium stearate, lactose, colloidal silicon  
20           dioxide, talc, and glyceryl behenate.

          The core can be prepared by any suitable tableting technique known to those skilled in the art. For example, the active ingredient may be admixed with the excipient(s) and advantageously formed into a tablet using a conventional tableting press. Alternatively, the core may be formulated as  
25           capsules, soft capsules, mini-tablets, granules, pellets, or spheronized crystals if desired, using conventional techniques.

          In order to achieve the site-specific pharmaceutical formulation, the core is coated with an intermediate layer which is a swellable polymeric coating layer. The swellable polymeric coating layer delays the release of

mesalamine for a predetermined period of time, which period of time is dependent upon the thickness of the swellable polymeric coating. In other words, the thicker the swellable polymeric coating, the longer it delays the release of the active ingredient from the core of the formulation. Thus, the appropriate site for the release of the active ingredient can be determined prior to the preparation of the formulation, and the formulation is designed by applying the appropriate thickness of swellable polymeric coating layer to achieve the desired time delay required to reach the predetermined site of delivery prior to release of the active ingredient.

The site-specific delivery of mesalamine using the instant formulation relies in part upon the fact that the residency time in the small intestine is relatively uniform from patient to patient. Typically, the transit time in the small intestine is  $3 \pm 1$  hours according to S. Davis et al., *Gut* 27:886 (1986). The formulation of the present invention relies upon the use of a control mechanism which can recognize the entry into the small intestine, and the use of polymer(s) or copolymer(s) which prevent the release of mesalamine from the core for the time needed to pass through the small intestine segment of the digestive tract.

The swellable polymeric coating layer comprises a hydrophilic gelling polymer or copolymer that swells on contact with gastro-intestinal juices to form a film surrounding the core. The swellable polymeric coating layer which surrounds the core protects the integrity of the core and prevents the release of mesalamine during the transit in the small intestine.

The swellable polymeric coating layer may be comprised of any suitable hydrophilic gelling polymer known to those skilled in the art. For example, suitable hydrophilic gelling polymers include but are not limited to cellulose polymers such as methylcellulose, carboxymethylcellulose, hydroxypropylcellulose, hydroxyethylcellulose, hydroxypropylmethylcellulose, and the like, vinyl polymers such as polyvinylpyrrolidone, polyvinyl alcohol, and the like, acrylic polymers and copolymers such as acrylic acid polymer,



-7-

methacrylic acid copolymers, ethyl acrylate-methyl methacrylate copolymers, and the like; natural or synthetic rubbers, poloxamers, polysaccharides, and mixtures thereof. Currently the preferred swellable polymeric coating layer comprises hydroxypropylmethylcellulose.

5 Hydroxypropylmethylcellulose is a polymer which is available in many grades, including types of different weight average molecular weight, extremely different viscosity, and different substitution grade. In one preferred embodiment, the swellable polymeric coating layer comprises a relatively low viscosity hydroxypropylmethylcellulose polymer having 1) a typical weight  
10 percent substitution corresponding to 29% methoxyl and 8% hydroxypropoxyl groups, and 2) a nominal viscosity of a 2% watery solution at 20°C ranging from 3 to 100 mPa.s, such as METHOCEL E5®, METHOCEL E50®, OR METHOCEL E100®, all available from Colorcon or PHARMACOAT 603® available from Seppic.

15 In addition to the foregoing, the swellable polymeric coating layer may also include additional excipients such as plasticizers, antisticking agents, and colorants. Specific examples of additional excipients include polyethylene glycol, polyvinylpyrrolidone, talc, magnesium stearate, glyceryl behenate, stearic acid, and titanium dioxide.

20 The swellable polymeric coating layer may be applied to the core using conventional film (or spray) coating techniques or double press coating. Preferably, the swellable polymeric coating layer is applied using the film coating techniques whereby the polymer is solubilized in an aqueous solution. Typically, the polymer used for film coating exhibits a viscosity ranging from  
25 about 3 to 100 mPa.s at 25°C in a 2% aqueous solution. Higher viscosity polymers can be applied using organic solutions or double press coating. Although some organic solvents may be employed in the film coating application of the swellable polymeric coating layer, the inclusion of organic solvents in the film coating solution is not required.

-8-

The solution of swellable polymer can be applied to the core by any means of film coating including but not limited to fluid bed, and pan coating. Preferably, the aqueous solution swellable polymer is sprayed on the core to form the swellable-polymeric coating layer.

5 The swellable polymer is applied to the core (preferably by film-coating) in order to build the desired thickness of the swellable polymeric coating layer. For example, in the embodiment wherein film coating is employed, the core is sprayed with the solution of polymer until the desired thickness of swellable polymeric coating layer is achieved. The desired thickness of the  
10 swellable polymeric coating layer is dependent upon the desired delivery site of the active ingredient. The thicker the swellable polymeric coating layer around the core, the longer the latency, or lag time prior to delivery of the mesalamine, and thus the farther through the gastro-intestinal tract the mesalamine will be delivered. Typically, the swellable polymeric coating layer is applied to a  
15 thickness sufficient to achieve a weight gain of between about 5 and about 200 percent, preferably between about 10 and about 100 percent as determined by solid substance. The weight ratio of the core:swellable polymeric coating layer is typically between about 20:1 and about 1:5, providing a thickness of swellable polymeric coating layer in excess of about 30  $\mu\text{m}$ , and up to about 3 mm.  
20 Preferably the ratio of core:swellable polymeric coating layer is between about 5:1 and about 1:3 inclusive, or a thickness of between about 100  $\mu\text{m}$  and about 1600  $\mu\text{m}$ .

The dissolution, disintegration, or erosion of the swellable polymeric coating layer at the desired site of delivery is triggered by the  
25 dissolution, erosion, or disintegration of the outer gastro-resistant layer. Once the predetermined delivery site is reached and the outer coating layer is dissolved, the swellable polymeric coating layer should be capable of relatively quick swelling and dissolution. The use of the low viscosity cellulose polymers for the preparation of the swellable polymeric coating layer according to the  
30 present invention provides a distinct advantage in this respect. The low viscosity

cellulose polymers which are used according to the present invention are characterized by a quicker solubilization, dissolution, or erosion time as compared to the high viscosity polymers. This relatively quick erosion time facilitates the maintenance of thinner levels of a thin swollen polymeric layer which is generated upon hydration. In other words, the erosion of the swollen polymeric coating layer proceeds uniformly. The achievement of a thin swollen layer permits a higher speed of release of mesalamine once the swellable polymeric coating layer has completely interacted and dissolved. The quick release of mesalamine provides a complete availability of the drug at the desired site of delivery.

The outer enteric coating layer of the instant formulation is a gastro-resistant coating layer, which permits the transit of the intact formulation through the stomach until the duodenum is reached. The outer coating layer overlies the swellable polymeric coating layer.

The outer enteric coating layer is comprised of conventional gastro-resistant polymers. For example, suitable gastro-resistant polymer for the outer enteric coating layer include acrylic/methacrylic copolymers, polyacrylates, polymethacrylates, acetate-phthalate cellulose, cellulose acetate terephthalate, cellulose acetate trimellitate, hydroxypropylmethylcellulose phthalate, or polyvinyl alcohol phthalate. Preferably, the outer enteric coating layer comprises acrylic/methacrylic copolymers such as those commercially available under the trade name EUDRAGIT®.

The outer enteric coating layer comprising the gastro-resistant polymer preserves the integrity of the formulation and inhibits the start of the swelling of the swellable polymeric coating layer during the gastric transit. The dissolution of the outer enteric coating layer is triggered by the presence of gastro-intestinal media having a pH above about 4.5, such as that present in the small intestine. Preferably, the dissolution of the outer enteric coating layer occurs when the formulation is subjected to a pH above about 5, and often above about 5.5. The dissolution of the outer enteric coating layer initiates the swelling

-10-

of the swellable polymeric coating layer. Thus, once the outer enteric coating layer is dissolved, the swelling, dissolution, and erosion of the swellable polymeric coating layer begins. The dissolution, erosion, or disintegration of the swellable polymeric coating layer inhibits the release of active ingredient from the core until the desired site of delivery is reached.

The outer enteric coating layer may be applied using any conventional coating techniques, including for example film coating techniques as described above. Preferably, the core coated with the swellable polymeric coating layer is further coated with the outer enteric coating layer using conventional spray coating techniques, wherein the coating process is carried out with a solution or suspension containing the gastro-resistant polymer which forms the outer enteric coating layer agent solubilized or suspended in a suitable solvent. Solvents useful for preparing the solution containing the outer coating agent include any pharmaceutically acceptable solvents capable of solubilizing the selected outer coating agent. A preferred solvent for preparing the solution containing the outer coating agent is water.

The site-specific dosage formulation of the present invention is suitable for oral administration and delivery in the colon. In order to achieve the site-specific delivery of mesalamine or other anti-inflammatory active ingredient, one need only administer the site specific formulation of the present invention to a subject in need of the a therapeutically effective dose of anti-inflammatory in the colon. Subjects in need of such treatment include humans, particularly humans suffering from ulcerative colitis or other inflammatory bowel disorders.

The formulation of the present invention provides a number of distinct advantages over conventional mesalamine formulations. As noted above, the formulation of the present invention utilizes often low viscosity cellulose polymers as the swellable polymeric coating layer, thus avoiding the necessity of including an organic solvent in the film coating process. Also, the low viscosity cellulose polymers permit the quick release of mesalamine at the pre-determined site. In addition, the formulation of the present invention provides the further

-11-

advantage that the release pattern of mesalamine after the dissolution of the outer enteric coating layer is not dependent upon or controlled by pH.

The following examples are provided to illustrate the present invention, and should not be construed as limiting thereof. In these examples,

5 "mg" means milligrams, "g" means grams, "mm" means millimeters, " $\mu$ m" means micrometers, "kp" means 9.807 Newton, "min." means minute(s), and " $^{\circ}$ C" means degrees Centigrade. All percentages are in percent by weight of the tablet unless otherwise indicated. Dissolution/disintegration tests are carried out according to the standard procedures set forth in the United States  
10 Pharmacopoeia for testing the dissolution/disintegration of tablets.

#### EXAMPLE 1

Tablet cores (20,000) containing 400 mg of mesalamine are prepared with the following composition:

	Mesalamine	400	mg
15	Lactose	57	mg
	Crospovidone	20	mg
	Talc	8	mg
	Glyceryl behenate	8	mg
	Polyvinylpyrrolidone	7	mg

20 Mesalamine is granulated with 10% solution of polyvinylpyrrolidone in order to obtain a homogeneous granulate. Thereafter, the wet mass is sieved and dried at 40 $^{\circ}$ C for 6 hours. The dried granules are calibrated through an adapted screen and then mixed with crospovidone, lactose, talc, and glycerol behenate in a ribbon mixer. The granular mixture is formed  
25 into tablet cores of 10mm in diameter, weighing 500 mg each using a rotary tablet press. The cores show a disintegration time lower than 5 minutes in water, a Schleuninger hardness higher than 10 kp and a friability lower than 0.1%.

-12-

The inner layer is applied onto the tablet cores in an automatic coating pan using the following solution:

Hydroxypropylmethycellulose (Methocel E50®)	7.5% w/w
PEG 6000	1.5% w/w
Water	91.0% w/w

Samples having a weight gain included from 10% to 50% are of the core weight are collected for analysis.

The results obtained are summarized in the following table:

	Weight ( $n=30 \pm S.D.$ )	Tablet height ( $n=30 \pm S.D.$ )	Layer Thickness (microns)
Uncoated Core	$507.6 \pm 44.5$	$6.595 \pm 0.051$	
+10%	$558.2 \pm 10.3$	$6.998 \pm 0.078$	200
+20%	$610.4 \pm 13.6$	$7.368 \pm 0.081$	385
+30%	$662.2 \pm 12.1$	$7.676 \pm 0.081$	540
+40%	$711.6 \pm 15.8$	$7.951 \pm 0.094$	680
+50%	$767.0 \pm 13.6$	$8.275 \pm 0.068$	840

Subsequently the outer layer is applied by continuous spraying of an aqueous gastroresistant suspension of CAP (Cellulose Acetate Phthalate), containing triacetin as plasticizer.

The disintegration test carried out on the finished tablets shows the absence of release at pH lower than 5 for at least 2 hours. Upon increasing the pH of the medium up to 7.5 the disintegration of the tablets occurs in  $256 \pm 10.8$  minutes.

-13-

**EXAMPLE 2**

Tablet cores containing 400 mg of mesalamine are prepared as described in the example 1, and press-coated with a mixture of hydroxypropylmethylcellulose (Methocel E50™, available from Colorcon), polyvinylpyrrolidone and polyethylenglycol, in the ratio 7:2:1, by means of a press-coating machine; to obtain a press-coated tablet having a weight of 800 mg.

The tablets are coated by a continuous spraying of a gastroresistant film of methacrylic ester aqueous suspension using an automatic pan.

**EXAMPLE 3**

The site-specific dosage-formulation of the present invention allows the release of mesalamine after the arrival of the formulation in the large bowel (ascending colon, transverse colon and descending colon). This phenomenon was observed by scintigraphic imaging in human volunteers orally dosed with tablet containing mesalamine formulated according to the present invention and labeled with <sup>153</sup>Samarium. This result suggests that this technology increases the topical therapeutic effect of mesalamine while reducing the systemic absorption. The reduced systemic absorption of mesalamine is well documented by the much lower blood peak concentration of mesalamine (about 10-fold lower than a conventional formulation) and percentage of urinary excretion of the metabolite n-acetylmесalamine (6% vs 19-30% of conventional formulation) which is a marker of systemic absorption.

The foregoing is illustrative of the present invention and is not to be construed as limiting thereof. The invention is defined by the following claims, with equivalents of the claims to be included therein.

-14-

**That Which Is Claimed Is:**

1. A pharmaceutical formulation for the site-specific delivery of mesalamine in the colon, said formulation comprising:
  - a) a core comprising mesalamine in an amount effective to produce a therapeutic anti-inflammatory effect;
  - 5        b) a swellable polymeric coating layer substantially surrounding said core, wherein said swellable polymeric coating layer inhibits the release of said mesalamine for a predetermined period of time dependent upon the thickness of said swellable polymeric coating layer, and wherein dissolution of said swellable polymeric coating layer is independent of pH to which said  
10        swellable polymeric coating layer is exposed; and
  - c) an outer enteric coating layer substantially surrounding said swellable polymeric coating layer, wherein said outer enteric coating layer dissolves upon exposure to pH greater than about 4.5, and wherein the  
15        dissolution of the outer enteric coating layer initiates the subsequent swelling of the swellable polymeric coating layer.
2. The pharmaceutical formulation according to Claim 1, wherein said core further comprises at least one pharmaceutically acceptable excipient.
3. The pharmaceutical formulation according to Claim 1, wherein said swellable polymeric coating layer comprises a hydrophilic gelling polymer selected from the group consisting of methylcellulose, carboxymethylcellulose, hydroxypropylcellulose, hydroxyethylcellulose, hydroxypropylmethylcellulose, polyvinylpyrrolidone, polyvinyl alcohol, acrylic acid polymer, methacrylic acid copolymers, ethyl acrylate-methyl methacrylate copolymers, natural rubbers, synthetic rubbers, poloxamers, polysaccharides, and mixtures thereof.



-15-

4. The pharmaceutical formulation according to Claim 1, wherein said swellable polymeric coating layer is sufficiently thick to achieve a weight gain based upon the weight of the core of between about 5 and 200 percent.

5. The pharmaceutical formulation according to Claim 1, wherein said swellable polymeric coating layer is not less than about 30  $\mu\text{m}$  thick.

6. The pharmaceutical formulation according to Claim 1, wherein said swellable polymeric coating layer is between about 30  $\mu\text{m}$  and about 3 mm thick.

7. The pharmaceutical formulation according to Claim 1, wherein said swellable polymeric coating layer comprising hydroxypropylmethylcellulose.

8. The pharmaceutical formulation according to Claim 1, wherein said enteric coating layer comprises a gastro-resistant polymer selected from the group consisting of acrylic/methacrylic copolymers, polyacrylates, polymethacrylates, acetate-phthalate cellulose, cellulose acetate terephthalate, cellulose acetate trimellitate, hydroxypropylmethylcellulose phthalate, or polyvinyl alcohol phthalate.

9. A method for achieving the site-specific delivery of mesalamine in the colon of a subject in need of such treatment, said method comprising orally administering to said subject, a site-specific dosage formulation comprising a) a core comprising mesalamine in an amount effective to produce a therapeutic anti-inflammatory effect, b) a swellable polymeric coating layer substantially surrounding said core, wherein said swellable polymeric coating

-16-

layer inhibits the release of mesalamine for a predetermined period of time dependent upon the thickness of said swellable polymeric coating layer, and wherein dissolution of said swellable polymeric coating layer is independent of pH to which said swellable polymeric coating layer is exposed; and c) an outer enteric coating layer substantially surrounding said swellable polymeric coating layer, wherein said outer enteric coating dissolves upon exposure to pH greater than about 4.5, and wherein the dissolution of said outer enteric coating layer initiates the swelling of said swellable polymeric coating layer.

10. The method according to Claim 9, wherein said core further comprises at least one pharmaceutically acceptable excipient.

11. The method according to Claim 9, wherein said swellable polymeric coating layer comprises hydrophilic polymer selected from the group consisting of methylcellulose, carboxymethylcellulose, hydroxypropylcellulose, hydroxyethylcellulose, hydroxypropylmethylcellulose, polyvinylpyrrolidone, polyvinyl alcohol, acrylic acid polymer, methacrylic acid copolymers, ethyl acrylate-methyl methacrylate copolymers, natural or synthetic rubbers, poloxamers, polysaccharides, and mixtures thereof.

12. The method according to Claim 9, wherein said swellable polymeric coating layer is sufficiently thick to achieve a weight gain based upon the weight of the core of between about 5 and 200 percent.

13. The method according to Claim 9, wherein said swellable polymeric coating layer is not less than about 30  $\mu\text{m}$  thick.

14. The method according to Claim 9, wherein said swellable polymeric coating layer is between about 30  $\mu\text{m}$  and about 3 mm thick.

-17-

15. The method according to Claim 9, wherein said swellable polymeric coating layer comprising hydroxypropylmethylcellulose.

16. The method according to Claim 9, wherein said enteric coating layer comprises a gastro-resistant polymer selected from the group consisting of acrylic/methacrylic copolymers, polyacrylates, polymethacrylates, acetate-phthalate cellulose, cellulose acetate terephthalate, cellulose acetate trimellitate, hydroxypropylmethylcellulose phthalate, or polyvinyl alcohol phthalate.



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 9/50, 31/60, 9/28</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 98/26767</b> <b>(43) International Publication Date:</b> 25 June 1998 (25.06.98)
<b>(21) International Application Number:</b> PCT/IB97/01652 <b>(22) International Filing Date:</b> 16 December 1997 (16.12.97) <b>(30) Priority Data:</b> 08/767,980 17 December 1996 (17.12.96) US <b>(71) Applicant (for all designated States except US):</b> POI.I INDUS- TRIA CHIMICA S.P.A. [IT/IT]; Piazza Agrippa, 1, I-20141 Milano (IT). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> Buseti, Cesare [IT/IT]; Via Signorini, 7, I-20149 Milano (IT). CRIMELLI, Tiziano [IT/IT]; Via Londonio, 1, I-20154 Milano (IT). OLGATI, Vincenzo [IT/IT]; Via Settembrini, 16, I-20010 Casorezzo (IT). <b>(74) Agents:</b> McCallum, William, Potter et al.; Cruikshank & Fairweather, 19 Royal Exchange Square, Glasgow G1 3AE (GB).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims</i> <i>and to be republished in the event of the receipt of amendments.</i>  <b>(88) Date of publication of the international search report:</b> 3 September 1998 (03.09.98)
<b>(54) Title:</b> SITE-SPECIFIC CONTROLLED RELEASE DOSAGE FORMULATION FOR MESALAMINE  <b>(57) Abstract</b> <p>Pharmaceutical formulation for the site-specific delivery of mesalamine in the colon is disclosed. The formulation includes (a) a core comprising mesalamine in an amount effective to produce a therapeutic anti-inflammatory effect; (b) a swellable polymeric coating layer substantially surrounding the core which inhibits the release of mesalamine for a predetermined period of time dependent upon the thickness of the swellable polymeric coating layer, and (c) an outer enteric coating layer substantially surrounding the swellable polymeric coating layer. Methods of achieving the site-specific delivery of mesalamine in the colon are also disclosed.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## INTERNATIONAL SEARCH REPORT

International Application No  
/IB 97/01652

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K9/50 A61K31/60 A61K9/28

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 2 245 492 A (ZAMBON SPA) 8 January 1992  see page 36 - page 37: example 16	1-4, 7-12, 15, 16
X	EP 0 673 645 A (HOFFMANN LA ROCHE) 27 September 1995  see page 6; example 1 see claim 1	1-3, 5, 8-11, 13-16
X	EP 0 671 167 A (TANABE SEIYAKU CO) 13 September 1995 see column 11 - column 12; example 4 see claims 1, 5, 7	1-16
	-/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## Special categories of cited documents

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"3" document member of the same patent family

Date of the actual completion of the international search

10 July 1998

Date of mailing of the international search report

21.07.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040. Tx 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Boulois, D

## INTERNATIONAL SEARCH REPORT

International Application No

IB 97/01652

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 453 001 A (GIULIANI SPA) 23 October 1991  see page 4; example 2	1-4, 7-12, 15, 16
X	WO 95 28963 A (WATANABE SHUNSUKE ; FUKUI MUNEKO (JP); KAWAI HITOSHI (JP); KATSUMA M) 2 November 1995 see the whole document	1-4; 8-12, 16
X, P	& EP 0 759 303 A (YAMANOUCHI PHARMACEUTICAL CO LTD) 26 February 1997 see page 14; example 4 see page 6, line 59 - page 7, line 32	1-4, 8-12, 16
X	EP 0 527 942 A (PERIO PROD LTD ; YISSUM RES DEV CO (US)) 24 February 1993 see page 2, line 1 - line 5 see page 3, line 6 - line 32 see page 4; example 1 see claims 1, 3, 8	1-16

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

/IB 97/01652

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 2245492 A	08-01-1992	IT 1244867 B	12-09-1994
		IT 1244037 B	28-06-1994
		AT 400295 B	27-11-1995
		AT 131091 A	15-04-1995
		AU 638583 B	01-07-1993
		AU 8019991 A	09-01-1992
		BE 1004882 A	16-02-1993
		CA 2044398 A	05-01-1992
		CH 683498 A	31-03-1994
		DE 4122039 A	09-01-1992
		DK 129591 A	05-01-1992
		ES 2036457 B	01-03-1994
		FI 913248 A	05-01-1992
		FR 2664163 A	10-01-1992
		GR 91100283 A, B	26-08-1992
		IE 61651 B	16-11-1994
		IL 98525 A	23-07-1996
		JP 6024961 A	01-02-1994
		LU 87964 A	03-03-1992
		NL 9101161 A	03-02-1992
		PT 98188 A	29-05-1992
		SE 9102072 A	05-01-1992
		RU 2012330 C	15-05-1994
		US 5310558 A	10-05-1994
		US 5445828 A	29-08-1995
		US 5629017 A	13-05-1997
EP 0673645 A	27-09-1995	US 5482718 A	09-01-1996
		AU 1488295 A	05-10-1995
		BR 9505785 A	05-03-1996
		CA 2142653 A	24-09-1995
		CN 1112417 A	29-11-1995
		JP 7277958 A	24-10-1995
		NZ 270737 A	26-11-1996
		ZA 9502169 A	26-09-1995
EP 0671167 A	13-09-1995	JP 7223970 A	22-08-1995
		CA 2142142 A	11-08-1995
		CN 1116926 A	21-02-1996
		US 5614220 A	25-03-1997



# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/IB 97/01652

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0453001 A	23-10-1991	IT 1246382 B	18-11-1994
		AT 144138 T	15-11-1996
		AU 654277 B	03-11-1994
		AU 7679891 A	11-11-1991
		CA 2035155 A	18-10-1991
		CA 2040471 A	18-10-1991
		DE 69122751 D	21-11-1996
		DE 69122751 T	13-02-1997
		DK 524989 T	18-11-1996
		WO 9116042 A	31-10-1991
		EP 0524989 A	03-02-1993
		ES 2093097 T	16-12-1996
		JP 4224517 A	13-08-1992
		PT 97370 A	31-01-1992
		SG 46584 A	20-02-1998
WO 9528963 A	02-11-1995	AU 689250 B	26-03-1998
		AU 2266895 A	16-11-1995
		CN 1146158 A	26-03-1997
		EP 0759303 A	26-02-1997
EP 0527942 A	24-02-1993	AU 660147 B	15-06-1995
		AU 8087791 A	27-11-1991
		DE 527942 T	03-03-1994
		ES 2048133 T	16-03-1994
		FI 924984 A	04-11-1992
		GR 94300025 T	31-05-1994
		HU 210497 B	28-04-1995
		HU 64210 A	28-12-1993
		IL 98087 A	14-11-1996
		WO 9116881 A	14-11-1991
		US 5525634 A	11-06-1996